

How much work can a quantum device extract from a heat engine?

D. Gelbwaser-Klimovsky,¹ R. Alicki,² and G. Kurizki¹

¹Weizmann Institute of Science, 76100 Rehovot, Israel

²Institute of Theoretical Physics and Astrophysics, University of Gdańsk

Work extraction from a heat engine in a cycle by a quantum mechanical device (quantum “piston”) is analyzed. The standard definition of work fails in the quantum domain. The correct extractable work and its efficiency bound are shown to crucially depend on the initial quantum state of the piston. The transient efficiency bound may exceed the standard Carnot bound, although it complies with the second law. Energy gain (e.g. in lasing) is shown to drastically differ from work gain.

Introduction Despite efforts to reconcile quantum mechanics and thermodynamics over the years[1–4], their compatibility is still an open foundational problem, with crucial bearing on the performance bounds of heat machines in the quantum domain [5–11]. The need to resolve this problem becomes apparent when we consider the widely accepted division of energy ΔE_S drawn by a system from the bath in a closed cycle into work, W , and heat, Q , in the quantum domain [9]

$$\Delta E_S = \oint \text{tr}(\rho_S \dot{H}_S) dt = -W + Q;$$

$$W = - \oint \text{tr}\{\rho_S \dot{H}_S dt\}; \quad Q = \oint \text{tr}\{\dot{\rho}_S H_S dt\}. \quad (1)$$

Here $\rho_S(t)$ is the reduced density matrix of the system and $\dot{H}_S$ is the time-derivative of the parametrically (*classically*) driven system Hamiltonian $H_S(t)$. In this convention, $W > 0$ implies that work is performed in a cycle by the system on the driving device (a “piston”) at the expense of part of the heat energy supplied by a bath. Ideally, the piston can extract all the work allowed by Eq. (1) within the fundamental Carnot efficiency bound[11]. The debate concerning the proper ensemble-averaging of work fluctuations[11] *does not affect* the validity of Eq. (1). Yet, what is the counterpart of Eq. (1) when we consider a *time-independent* Hamiltonian H_{tot} governing a closed complex, whose ingredients (the baths, the system and the piston) are fully quantized? Since $\dot{H}_{tot} = 0$, does this imply that work *cannot* be extracted at all? And if it can, do quantum correlations or coherence in the complex play any role in determining the work extraction?

Here we attempt to elucidate these widely open fundamental questions, and thereby the formidable issue of the compatibility of quantum mechanics and thermodynamics, by resorting to the seldom used notion of the “*passivity*” of a mixed quantum state[12]: the definition of work extraction as the difference between the mean energy in this state and in the corresponding passive state (the state of *minimal* mean energy at the same entropy). We show that it is necessary to use the passivity definition of work in quantized closed complexes to avoid violation of the second law, with which the *standard* (canonical) definition [5–9] confronts us in such scenarios. We

therefore use passivity to infer the bound on work extraction by a quantum piston and investigate the rapport of this bound with Carnot efficiency. The investigation shows that this bound may drastically differ from the conventional Carnot bound[13]: it involves *effective temperature and entropy* that strongly depend on the initial quantum state and dynamics of the piston. Strikingly, the transient efficiency for an initial coherent state may exceed the standard Carnot bound, thus revealing the ability of the quantum piston to act as a *negentropy resource*. The general results are supported by an analysis of a generic, fully quantized model, which may describe maser/laser action of a “piston” (signal) mode in a cavity [5, 7, 14]. This model reveals *the principal discrepancy between work and energy gain* of the piston mode.

Canonical bound We consider a system (S) coupled to cold (C) and hot (H) baths and to a piston (P), all fully quantized. Since the standard Eq. (1) does not apply when the Hamiltonian is time-independent, we may instead invoke the first law of thermodynamics (energy conservation) for the total (closed) complex of the system (S), piston (P), cold (C) and hot (H) baths, assuming the engine operates at the steady-state of the system, so that $\frac{d\langle H_S \rangle}{dt} = 0$. The first law then yields the standard (*canonical*) expression for power generation by the engine [9] $\mathcal{P}_{can} \equiv \frac{d\langle H_P \rangle}{dt} = (J_C + J_H)$, where the energy flow between the two baths and S is given by the respective heat currents, J_C , and J_H [6, 9, 15]. The *canonical* efficiency bound is then the ratio of the output power $\mathcal{P}_{can}$ to the heat current input J_H

$$\eta_{can} = \frac{\mathcal{P}_{can}}{J_H} = \frac{J_C + J_H}{J_H} \quad (2)$$

However, if the canonical expression (2) indeed pertains to the rate of work and the efficiency, whereas J_C and J_H are calculated while allowing for the state of a quantum piston (see below), then the Kelvin formulation of the second law, whereby work cannot be extracted in a cycle by a single-bath engine, may be violated (Fig. 1a).

In an attempt to remove such inconsistencies we may invoke Spohn’s theorem[16], which is a statement of the second law in the Markovian approximation[1]: $\dot{S}_P \geq \frac{J_C}{T_C} + \frac{J_H}{T_H}$. Here the l.h.s is the entropy production-rate of




FIG. 1. (a) Work extraction/(initial work stored in the piston) from a *single bath*, $T_C = T_H$, for a piston initially in a coherent state (with amplitude $\alpha(0) = 1/5$), calculated for the model of Eq.(6). The canonical bound (Eq. (2)) permits work to be extracted from a single bath in contrast to the passivity bound (Eq. (4)) in keeping with the second law. The cycle duration is $\frac{2\pi}{\nu}$. Insets: lower–piston mode drives quantum dots embedded between two materials at different temperatures. Upper– a lasing piston mode in a cavity setup. (b) Transient efficiency bound and work extraction (initial work) according to the passivity bound Eq. (4) for $\frac{T_C}{T_H} = 1/50$ and a coherent state with initial energy $\nu\alpha(0)^2 = 0.98$

the piston, which is commonly ignored [6–10], but may be important for a quantum piston. Under this inequality we find that the canonical efficiency bound (2) for a cycle, at given bath temperatures T_C and T_H , satisfies $\eta_{can} \leq 1 - \frac{T_C}{T_H} + \frac{T_C}{J_H} \dot{S}_P$. This inequality implies the possibility to violate both the Carnot bound for $T_H > T_C$ and Kelvin’s prohibition on work extraction from a single bath ($T_C = T_H$), on account of the term $\frac{T_C}{J_H} \dot{S}_P$, which is positive-definite (in the Markovian approximation)[1]. Yet this term describes *heat exchange* between the piston and the engine, rather than work extraction. We conclude that the *canonical definition of output power is inappropriate for the work-extraction bound* when the entropy production-rate of the quantum piston plays a role, as it fails to distinguish between work and heat pro-

duction.

Passivity bound The starting point of our alternative analysis is the definition of work based on the notion of passivity[12], whereby the work extractable by the piston (P) in a given state is $W_P = \langle H_P(\rho_P) \rangle - \langle H_P(\rho'_P) \rangle_{min}$ where H_P is the *time-independent* piston Hamiltonian, ρ_P denotes the density matrix of the piston (obtained upon tracing out the system S and the baths B_C and B_H), and $\rho'_P = U\rho_P U^\dagger$ is its counterpart following a unitary transformation that minimizes the second term: i.e. we have to find a unitary such that W_P is maximized, by minimizing $\langle H_P(\rho'_P) \rangle$. It is easy to find the passivity bound on work extraction in a cycle:

$$(W_P)_{Max} = \langle H_P(\rho_P) \rangle - \langle H_P(\rho'_P) \rangle_{Gibbs};$$

$$(\rho'_P)_{Gibbs} = Z^{-1} e^{-\frac{H_P}{T_P}} \quad (3)$$

where we have used the property of the Gibbs (equilibrium) state that minimizes the energy for a *fixed* entropy $\mathcal{S}_P$ (an invariant under unitary transformations). The Gibbs form yields the *effective* temperature T_P of the piston from its invariant entropy $\mathcal{S}_P$. We stress that it is by no means guaranteed that ρ_P can be transformed to a Gibbs state by a unitary process, so that the bound might not be tight. However, applying entangling unitaries to an ensemble of “n” such pistons one can reach asymptotically this bound [17].

The variation of Eq. (3) (the passivity bound) under energy exchange between S, P, B_H and B_C yields $(\mathcal{P}_{pas})_{Max} = \frac{d(W_P)_{Max}}{dt} = \frac{d\langle H_P \rangle}{dt} - T_P \dot{S}_P$, where we have identified the power extraction with the rate of change of work by the piston. The last term on the r.h.s, involving the entropy production-rate of the piston, is the difference between the power extraction by the piston according to the passivity definition and the canonical expression. The passivity bound on the power yields the efficiency bound

$$(\eta_{pas})_{Max} = \frac{J_C + J_H - T_P \dot{S}_P}{J_H}. \quad (4)$$

This bound can be checked not to violate Kelvin’s prohibition on single-bath work extraction as opposed to Eq. (2) (Fig. 1a). However, the compliance of (4) with the Carnot bound is not obvious. Using Spohn’s theorem [16] as before, now yields, instead of the canonical bound $\eta_{pas} \leq 1 - \frac{T_C}{T_H} + \frac{(T_C - T_P) \dot{S}_P}{J_H}$. This inequality agrees with the Carnot bound if $T_C \geq T_P$. Yet what if this is not the case? For $T_P \leq T_C$, the Spohn theorem [16] and the convention [15] $J_C \leq 0$ imply the inequality $\dot{S}_P \geq \frac{J_C}{T_C} + \frac{J_H}{T_H} \geq \frac{J_C}{T_P} + \frac{J_H}{T_H}$, whence

$$\eta_{pas}(T_P \leq T_C) \leq 1 + \frac{J_H - T_P(\frac{J_H}{T_P} + \frac{J_H}{T_H})}{J_H} = 1 - \frac{T_P}{T_H} \quad (5)$$

This bound implies that *quantum engine effectively operates between three different temperatures: T_H , T_C and T_P* . Its maximum efficiency is the Carnot bound $1 - \frac{T_C}{T_H}$ for $T_C \leq T_P \leq T_H$, but for $T_P \leq T_C$ it is $1 - \frac{T_P}{T_H}$ i.e. it *exceeds* the standard Carnot bound $1 - \frac{T_C}{T_H}$ (Fig. 1b). Yet, it *complies* with Spohn's theorem and hence with the *second law*. This remarkable result prompts the questions: what determines the discrepancy between this passivity bound, the canonical bound and the standard Carnot bound? What determines the *actual* work extraction (rather than its bound) by a quantum piston?

Model analysis The fundamental questions raised above call for an analysis of the *simplest (minimal)* model of a quantum heat engine (QHE), to date: a harmonic-oscillator (piston) that acts *dispersively* on a *single-qubit* system (i.e. without changing the level populations), while the system is permanently coupled to *spectrally-separated* hot (H) and cold (C) baths at temperatures T_H and T_C , respectively. The parametrically (classically) piston-modulated version of this model has been studied by us [15]. We wish to understand: How different is its quantum counterpart?

To this end, we consider the quantized model of dispersive S-P coupling (Suppl. Info.) :

$$H_{S+P} = \frac{1}{2}\omega_0\sigma_Z + \nu a^\dagger a + \frac{g_p}{2}(a^\dagger + a)\sigma_Z \quad (6)$$

where the first term is the two-level system (TLS) energy, the second term is the quantized-piston energy and the last term is the weak S-P interaction at a rate g_p that couples the sum of the creation and annihilation operators of P to the σ_Z Pauli operator of S. The S-B coupling has the standard spin-boson form $H_{SB} = \sigma_X(B_H + B_C)$.

We wish to analyze the piston evolution in this model. We describe its bath-induced dynamics by the (interaction picture) Lindblad-Gorini-Kossakowski-Sudarshan (LGKS) generator obtained in the Davies weak coupling limit procedure [2] (Suppl. Info.) $\mathcal{L} = \sum_j \sum_q \mathcal{L}_{j,q}$ where $j = H, C$ labels baths and $q = 0, \pm 1$ labels relevant Bohr frequencies $\omega_0 + q\nu$ of $S + P$. According to [9] the time-dependent currents read $J_j = \sum_q \text{Tr}(H_{S+P} \mathcal{L}_{j,q} \rho_{S+P}(t))$ [15]. Then using a “dressing transformation” (Suppl. Info) the (dressed) TLS reaches a steady state while the piston still evolves, thus we can eliminate the TLS variables. This yields the final description in terms of (interaction picture) LGKS master equation for the (dressed) harmonic oscillator- the piston. Having the time dependent reduced density matrix of the piston $\rho_P(t)$ we can compute its entropy $S(\rho_P)$ and its effective temperature T_P .

For further insight, we cast the Liouville (master) equation for the piston into a physically transparent form in the Glauber-Sudarshan P-representation [14, 18]. The quasiprobability distribution in the coherent-state basis, $\rho_P = \int d^2\alpha \mathbf{P}(\alpha) |\alpha\rangle\langle\alpha|$, then obeys the Fokker-Planck equation




FIG. 2. Main panels: Non-passivity of $\mathbf{P}(\alpha)$ for an initial (a) Fock state, (b) coherent state, and the corresponding work-extraction evolution states: 1) pointed (after 2.310^{10} cycles); 2) dashed (after 3.1×10^{10} cycles); 3) solid (after 7.8×10^{10} cycles). Below: the corresponding phase-plane distributions at (c) (stage 1) and (d) (stage 3).

$$\frac{d\mathbf{P}}{dt} = \frac{\gamma}{2} \left(\frac{\partial}{\partial \alpha} \alpha + \frac{\partial}{\partial \alpha^*} \alpha^* \right) \mathbf{P} + D \frac{\partial^2 \mathbf{P}}{\partial \alpha \partial \alpha^*}, \quad (7)$$

$$\gamma = \frac{g_p^2}{\nu^2} \left((G(\nu_+) - G(\nu_-)) \rho_{ee} + (G(-\nu_-) - G(-\nu_+)) \rho_{gg} \right) \quad (8)$$

$$D = \frac{g_p^2}{\nu^2} ((G(\nu_-) \rho_{ee} + G(-\nu_+) \rho_{gg})) \quad (9)$$

where $\nu_{\pm} = \omega_0 \pm \nu$, γ is the drift (loss or gain) coefficient, D is the diffusion coefficient and $G(\omega) = \sum_{j=H,C} G^j(\omega)$ is the sum of the cold- and hot-baths response spectra. The passivity expression for work then assumes the form

$$W_P = \nu \int d^2\alpha |\alpha|^2 (\mathbf{P}(\alpha) - \mathbf{P}'(\alpha)) \quad (10)$$

$\mathbf{P}'(\alpha)$ being the minimal-energy unitarily-transformed distribution. As in (3), the second term represents the contribution of the passive state. Hence, Eq. (10) expresses work extraction as *non-passivity*. This stands in stark contrast to the piston *energy gain or loss*:

$$\langle H_P(t) \rangle = \nu \left(\frac{D}{\gamma} (1 - e^{-\gamma t}) + e^{-\gamma t} \langle \alpha^2(0) \rangle \right) \quad (11)$$

which is independent of the passivity of the initial state.

How much work is extracted from a given initial state? The Fokker-Planck equation (7) allows us to analytically calculate the work in a cycle from (10) for a variety of initial states of the piston, if we can find the energy-minimizing unitary transformation $\mathbf{P}(\alpha) \rightarrow \mathbf{P}'(\alpha)$ (Fig. 2):

(i) For an initial *coherent state* $\alpha(0)$, work extraction, described by a unitary displacement of the distribution to $\alpha' = 0$ yields, $W_P = \nu \alpha^2(0) e^{-\gamma t}$ for any $\alpha(0)$. This work *exponentially increases for negative γ* . The quasiclassical limit retrieves the canonical expression (2) for power: $\lim_{\langle \alpha(0) \rangle^2 \rightarrow \infty} \mathcal{P}_{Passive} \rightarrow \frac{\langle dH_P \rangle}{dt}$.

(ii) “*Incoherent*” state, e.g. a phase-uniform distribution of coherent states with the same $|\alpha(0)|$, $\mathbf{P}(\alpha, \alpha^*) = \int_0^{2\pi} d\theta \delta^2(\alpha - \alpha(0))$, yields work extraction only in the *quasiclassical limit* $\alpha(0)^2 \gg 1$.

(iii) A *thermal state* will keep its character and not extract work, $W_P < 0$, which implies *passivity at any time*.

(iv) A pure *initial Fock state* $|n(0)\rangle$ can at most deliver the work $\nu n(0)$ initially stored in it, but contrary to the coherent state, *it cannot extract more work* from the engine.

We may thus conclude that the extracted work strongly depends on the initial state of the piston, particularly its *coherence*. The question is: what determines whether a state is suited for work extraction or not? To answer this question in general terms, we look for criteria for sustainable (long-time) non-passivity. This requires *non-monotonic decrease* of $\mathbf{P}(\alpha)$ with the energy $\nu|\alpha|^2$ at long times. This requirement may be formulated in terms of the derivative with respect to $r > 0$, where $\alpha = r e^{i\theta}$

$$\frac{\partial \mathbf{P}(\alpha, t)}{\partial r} > 0 \quad (12)$$

The following considerations emerge from the analysis of Eq. (12) (Suppl. Info):

(a) An initially passive (e.g., thermal) state, will remain so at later times, with no work extraction at any time, since Eq. (12) cannot be satisfied. (b) *Only $\gamma < 0$* can yield sustainable work, since for $\gamma > 0$ the distribution always becomes thermal distribution.

(c) The main contribution to Eq. (12) comes from $|\alpha|$ close to the center of the distribution. If the center stays at the origin as in a Fock state, the distribution will eventually become passive and terminate work extraction. By contrast, a coherent state, whose distribution is centered at $|\alpha| = |\alpha(0)| e^{-\frac{\gamma t}{2}}$ retains the nonpassivity, so that work extraction is sustainable at all times.

(d) For a piston in an initial passive or Fock state, *the engine needs an “ignition”, which can be achieved by coherence injection* expressed by $|P_{\text{ignition}}\rangle = D(\alpha)|P(0)\rangle$, where $D(\alpha)$ is the displacement operator and $|P(0)\rangle$ is the initial (here pure) state.

Exceeding the Carnot bound Finally, our analysis of the passivity bound (5) allows the engine to surpass *the standard Carnot bound* when the piston is initially in a coherent state (Fig 1b). This is allowed (by Eq. (4)) since the coherent-state effective temperature $T_P(t)$ then rises with time as

$$\frac{1}{T_P(t)} = \frac{1}{\nu} \text{Log} \left(\frac{1 + Dt}{Dt} \right). \quad (13)$$

Discussion The present analysis allows to understand work extraction in the quantum domain as *deviation from a passive state*. The compelling argument in favor of this seldom-used definition is that it avoids inconsistency with Kelvin’s second law, as opposed to the standard (canonical) definition. The passivity efficiency bound (Eqs. (4), (5)) assumes the form of a Carnot bound, if the time-dependent effective temperature $T_P(t)$ of the piston is introduced, yet for $T_P < T_C$ (Eq. (13)) it may surpass the standard Carnot bound. This striking transient effect shows that quantum coherence may provide *negentropy* beyond the standard (classical) limit of work extraction.

A possible experimental setup for demonstrating and further studying the effects discussed here may involve an ensemble of N two-level systems (TLS) realized by adsorbates or defects deposited at an interface between two solids at different temperatures, T_C and T_H . The solid structure also serves as a resonator for a radiation “piston” mode that is off-resonantly coupled to the TLS (Fig. 1-Inset). The hotter solid (at T_H) is the energy source for the engine that transforms phonons (via dipole coupling to the TLS) into the piston-mode excitation. A thermal-radiation (microwave) cavity with a piston mode is another experimental candidate (Fig 1-Inset). The energy-gain of the piston mode (Eq. (11)) is necessary for work extraction, but the latter may be *drastically different* from energy gain (Eqs. (10),(12)) depending on the initial state.

To conclude, the present results open a new vista into the obscure interface between engine thermodynamics and quantum dynamics. Hitherto unfamiliar quantum effects have been shown to affect the engine performance, particularly in the low-excitation (few quanta) limit.

The support of ISF, DIP, Grant No. NN 202208238 of MNiSW (RA) and CONACYT is acknowledged.

-
- [1] G. Lindblad, *Non-Equilibrium Entropy and Irreversibility* (D. Reidel, Holland, 1983).
 [2] E. Davies, Commun. Math. Phys. **39**, 91 (1974);
 G. Lindblad, Commun. Math. Phys. **40**, 147 (1975);

- V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, *Journal of Mathematical Physics* **17**, 821 (1976); R. Alicki and K. Lendi, *Quantum Dynamical Semigroups and Applications* (Springer, Berlin, Germany, 2007).
- [3] J. v. Neumann, *Proceedings of the National Academy of Sciences* **18**, 70 (1932); N. Linden, S. Popescu, A. J. Short, and A. Winter, *Phys. Rev. E* **79**, 061103 (2009); S. Goldstein, J. L. Lebowitz, C. Mastrodonato, R. Tumulka, and N. Zangh, *Proc. Royal Soc. A* **466**, 3203 (2010); M. Rigol and M. Srednicki, *Phys. Rev. Lett.* **108**, 110601 (2012).
- [4] J. Gemmer, M. Michel, and G. Mahler, *Quantum Thermodynamics* (Springer, Berlin, Germany, 2010).
- [5] J. E. Geusic, E. O. Schulz-DuBios, and H. E. D. Scovil, *Phys. Rev.* **156**, 343 (1967); G. W. Ford and R. F. O'Connell, *Phys. Rev. Lett.* **96**, 020402 (2006); G. W. Ford, J. T. Lewis, and R. F. O'Connell, *Phys. Rev. Lett.* **55**, 2273 (1985).
- [6] E. Geva, *J. Mod. Opt.* **49**, 635 (2002); H. T. Quan, Y.-x. Liu, C. P. Sun, and F. Nori, *Phys. Rev. E* **76**, 031105 (2007); C. Van den Broeck, *Phys. Rev. E* **95**, 190602 (2005); B. Lin and J. Chen, *ibid.* **67**, 046105 (2003); M. Esposito, *Phys. Rev. Lett.* **102**, 130602 (2009); T. D. Kieu, *ibid.* **93**, 140403 (2004); N. Linden, S. Popescu, and P. Skrzypczyk, *ibid.* **105**, 130401 (2010).
- [7] M. O. Scully, K. R. Chapin, K. E. Dorfman, M. B. Kim, and A. Svidzinsky, *Proceedings of the National Academy of Sciences* **108**, 15097 (2011).
- [8] V. Blickle and C. Bechinger, *Nature Physics* **7**, 1 (2011).
- [9] R. Alicki, *Journal of Physics A* **12**, 103 (1979).
- [10] A. E. Allahverdyan and T. M. Nieuwenhuizen, *Phys. Rev. E* **71**, 046107 (2005).
- [11] P. Talkner, E. Lutz, and P. Hänggi, *Phys. Rev. E* **75**, 050102(R) (2007); M. Campisi, P. Talkner, and P. Hänggi, *Phys. Rev. Lett.* **102**, 210401 (2009); M. Campisi, P. Hänggi, and P. Talkner, *Rev. Mod. Phys.* **83**, 771 (2011); C. Jarzynski, *Comptes Rendus Physique* **8**, 495 (2007).
- [12] A. Lenard, *Journal of Statistical Physics* **19**, 575 (1978); W. Pusz and S. Woronowicz, *Communications in Mathematical Physics* **58**, 273 (1978).
- [13] H. B. Callen, *Thermodynamics and an Introduction to Thermostatistics* (John Wiley & Son, Singapore, 1985).
- [14] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, UK, 1997).
- [15] D. Gelbwaser-Klimovsky, R. Alicki, and G. Kurizki, *Phys. Rev. E* (in press), arXiv:1209.1190v2 [quant-ph]; M. Kolář, D. Gelbwaser-Klimovsky, R. Alicki, and G. Kurizki, *Phys. Rev. Lett.* **109**, 090601 (2012).
- [16] H. Spohn, *J. Math. Phys.* **19**, 1227 (1978).
- [17] R. Alicki and M. Fannes, arXiv:1211.1209v2 [quant-ph] (2012).
- [18] H. Carmichael, *Statistical methods in quantum optics* (Springer, 1999); W. Schleich, *Quantum optics in phase space* (Wiley-VCH, 2001).

SUPPLEMENTARY INFORMATION

Dynamics in the fully quantized system-bath-piston complex

The S+P Hamiltonian reads

$$H_{S+P} = \frac{1}{2}\omega_0\sigma_Z + \frac{g_p}{2}(a + a^\dagger)\sigma_Z + \nu a^\dagger a \quad (S1)$$

This system is weakly coupled to “cold” and “hot” heat baths by the following interaction Hamiltonian

$$H_{int} = \sigma_X \otimes (B_C + B_H). \quad (S2)$$

The analysis of this model is simplified by using a new set of canonical operators $b, b^\dagger$ and Pauli matrices obtained by a unitary *dressing transformation* that diagonalizes the Hamiltonian

$$a \mapsto b = U^\dagger a U, \quad \sigma_k \mapsto \tilde{\sigma}_k = U^\dagger \sigma_k U, \quad U = e^{\frac{g_p}{2\nu}(a^\dagger - a)\sigma_z}. \quad (S3)$$

Under this transformation the Hamiltonian of S+P (S1) can be written as

$$\tilde{H}_{S+P} = \frac{1}{2}\omega_0\sigma^3 + \nu b^\dagger b - \left(\frac{g_p}{2}\right)^2 \frac{1}{\nu} \quad (S4)$$

and the dressing transformation can be expressed in terms of new variables

$$U = e^{\frac{g_p}{2\nu}U^\dagger(a^\dagger - a)\sigma_z}U = e^{\frac{g_p}{2\nu}(b^\dagger - b)\sigma_z}. \quad (S5)$$

The Pauli matrix σ_X which appears in the interaction Hamiltonian (S2) is given in terms of new dynamical variables as

$$\sigma_X = \tilde{\sigma}_+ e^{\frac{g_p}{\nu}(b^\dagger - b)} + e^{-\frac{g_p}{\nu}(b^\dagger - b)} \tilde{\sigma}_- \quad (S6)$$

The Heisenberg picture Fourier decomposition of σ_X within the lowest order approximation with respect to the small parameter g_p/Ω , can be obtained from

$$\begin{aligned} \sigma_+(t) &= e^{iHt}\sigma_+e^{-iHt} = e^{i\omega_0 t}\tilde{\sigma}_+e^{\frac{g_p}{\nu}(b^\dagger e^{i\nu t} - b e^{-i\nu t})} \approx \\ &\tilde{\sigma}_+e^{i\omega_0 t} + \frac{g_p}{\nu}(S_1^\dagger e^{i(\omega_0 + \nu)t} - S_{-1}^\dagger e^{i(\omega_0 - \nu)t}) \end{aligned} \quad (S7)$$

where

$$S_1^\dagger = \tilde{\sigma}_+ b^\dagger, \quad S_{-1}^\dagger = \tilde{\sigma}_+ b \quad (S8)$$

The approximation made in (S7) is valid for piston excitation numbers such that

$$\frac{g_p}{\nu}\sqrt{\langle b^\dagger b \rangle} \ll 1, \quad (S9)$$

The derivation of the coarse-grained master equation leads to the following (Interaction-picture) equation of motion for the reduced density matrix of S+P, ρ_{S+P}

$$\frac{d\rho_{S+P}(t)}{dt} = \mathcal{L}\rho_{S+P}(t) \quad (S10)$$

where

$$\mathcal{L} = \sum_{q=0,\pm 1} (\mathcal{L}_q^C + \mathcal{L}_q^H) \quad (\text{S11})$$

and the generators $\mathcal{L}_q^j$ are of the standard LGKS form

$$\mathcal{L}_0^j \rho_{S+P} = \frac{1}{2} \left\{ G^j(\omega_0) ([\tilde{\sigma}_- \rho_{S+P}, \tilde{\sigma}_+] + [\tilde{\sigma}_-, \rho_{S+P} \tilde{\sigma}_+]) + G^j(-\omega_0) ([\tilde{\sigma}_+ \rho_{S+P}, \tilde{\sigma}_-] + [\tilde{\sigma}_+, \rho_{S+P} \tilde{\sigma}_-]) \right\} \quad (\text{S12})$$

$$\mathcal{L}_q^j \rho_{S+P} = \frac{g^2}{2\nu^2} \left\{ G^j(\omega_0 + q\nu) ([S_q \rho_{S+P}, S_q^\dagger] + [S_q, \rho_{S+P} S_q^\dagger]) + G^j(-\omega_0 - q\nu) ([S_q^\dagger \rho_{S+P}, S_q] + [S_q^\dagger, \rho_{S+P} S_q]) \right\}, \quad q = \pm 1. \quad (\text{S13})$$

The effects of the baths are here described by the Fourier transforms of the autocorrelation functions ($j = C, H$)

$$G^j(\omega) = \int_{-\infty}^{+\infty} e^{i\omega t} \langle B^j(t) B^j \rangle_{T_j} dt = e^{\omega/T_j} G^j(-\omega), \quad (\text{S14})$$

The transition operators $S_{\pm 1}$ describe the processes of relaxation of the TLS accompanied by the excitation or deexcitation of the piston, while $S_{\pm 1}^\dagger$ describe their time-reversed counterparts.

Evolution in the Fock-state basis

The partially steady-state TLS populations in ρ_{S+P} are found to be related to the piston matrix elements $\rho_{nn'}$, n and n' being Fock states of P, by

$$(\rho_{nn'}^{ee})_{S+P} = \rho_{nn'} \frac{G(-\omega_0)}{G(-\omega_0) + G(-\omega_0)}; \quad (\rho_{nn'}^{gg})_{S+P} = \rho_{nn'} \frac{G(\omega_0)}{G(-\omega_0) + G(-\omega_0)}; \quad \rho_{nn'} = \rho_{nn'}^{gg} + \rho_{nn'}^{ee}. \quad (\text{S15})$$

Once the fast dynamics of S has reached steady state, it can be eliminated and effective equations can then be written for the *slow* degrees of freedom of P. In particular, the diagonal elements $\rho_{nn'}$, which are *decoupled* from the off-diagonal ones $\rho_{nn'}$, obey the form that is well known from quantum laser/maser theory [14]

$$\dot{\rho}_{nn} = r_{n+1}^+ \rho_{n+1,n+1} + r_{n-1}^- \rho_{n-1,n-1} - (r_n^+ + r_n^-) \rho_{n,n} \quad (\text{S16})$$

where r_n^+ and r_n^- denote the transition rates from $|n\rangle$ to $|n-1\rangle$ and vice versa, respectively. The specific features of our model are embodied by the dependence of $r_n^\pm$ on

the combined H and C bath spectra at the TLS resonance ω_0 , the sidebands $\omega_0 \pm \nu$ and their negative counterparts:

$$r_n^\pm = r^\pm n, \quad r^\pm = \left(\frac{g_p}{\nu}\right)^2 \frac{G(\nu_\pm)G(-\omega_0) + G(-\nu_\mp)G(\omega_0)}{G(\omega_0) + G(-\omega_0)} \quad (\text{S17})$$

The Glauber-Sudarshan P-representation is advantageous compared to the Fock-state basis, in that the P-representation offers unambiguous criteria for work extraction (see below). It may appear that initial quantum states with a *passive* (*monotonically-decreasing*) *distribution of Fock states populations*, $\rho_{nn} > \rho_{n+1n+1}$, are *unable to extract work*, but this may not be true, since an initial coherent (Glauber) state with $\langle n \rangle = 1$, $\rho_P = \sum_{i,j} \frac{|i><j|}{e^{\sqrt{i!j!}}}$ has a passive distribution of Fock-state populations, and *yet yields work*. On the other hand, the *same* Fock-state population distribution with random phases, $\rho_P = \sum_n \frac{|n><n|}{e^{(n!)}}$, i.e., without quantum coherence, yields no work. For a passive distribution of Fock-state populations, quantum coherence in the Fock basis $\rho_{nn'} \neq 0$ is a *necessary but insufficient* condition for work extraction, as we have checked numerically.

Work-extraction condition (Eq. 12) in the P-representation

The evolution of any initial distribution is described by [18]

$$P(re^{i\theta}, t) = \int r_0 dr_0 d\theta_0 \frac{k e^{-k|(re^{i\theta} - r_0 e^{-\gamma t/2} e^{i\theta_0})|^2}}{\pi} P(r_0 e^{i\theta_0}) \quad (\text{S18})$$

where $\alpha = re^{i\theta}$ and $\alpha_0 = r_0 e^{i\theta_0}$ where $k = \frac{\gamma}{D(1-e^{-\gamma t})}$. We are interested in its radial derivative

$$\frac{\partial P(re^{i\theta}, t)}{\partial r} = -2 \int dr_0 d\theta_0 (r - r_0 e^{-\gamma t/2} \cos(\theta - \theta_0)) r_0 \times \frac{k^2 e^{-k|(re^{i\theta} - r_0 e^{-\gamma t/2} e^{i\theta_0})|^2}}{\pi} P(r_0 e^{i\theta_0})$$

a) Let us assume an initial passive distribution, i.e., isotropic and $\frac{\partial P(r_0)}{\partial r_0} < 0$.

$$\frac{\partial P(re^{i\theta}, t)}{\partial r} = -2 \int dr_0 d\theta_0 (r - r_0 e^{-\gamma t/2} \cos(\theta - \theta_0)) r_0 \times \frac{k^2 e^{-k|(re^{i\theta} - r_0 e^{-\gamma t/2} e^{i\theta_0})|^2}}{\pi} P(r_0)$$

Integration over angle θ_0 yields

$$\frac{\partial P(re^{i\theta}, t)}{\partial r} = -4 \int \left(dr_0 (r I_0(2krr_0 e^{-\frac{\gamma t}{2}}) + r_0 e^{-\gamma t/2} I_1(-2krr_0 e^{-\frac{\gamma t}{2}})) r_0 k^2 e^{-k(r^2 + r_0^2 e^{-\gamma t})} P(r_0) \right)$$

where I_m is the modified Bessel function.

Integrating the r.h.s by parts yields

$$\begin{aligned} \frac{\partial P(re^{i\theta}, t)}{\partial r} = & \int dr_0 r 2r_0^2 \tilde{F}_1(2krr_0 e^{-\frac{\gamma t}{2}}) \frac{\partial k^2 e^{-k(r^2 + r_0^2 e^{-\gamma t})} P(r_0)}{\partial r_0} + \\ & \int dr_0 \frac{4r_0^2}{3} e^{-\gamma t/2} \frac{\partial I_1(-2krr_0 e^{-\frac{\gamma t}{2}}) k^2 e^{-k(r^2 + r_0^2 e^{-\gamma t})} P(r_0)}{\partial r_0} \end{aligned}$$

where ${}_0\tilde{F}_1(z)$ is the Regularized Hypergeometric Function which is positive for positive arguments. If $\frac{\partial P(r_0)}{\partial r_0} < 0$, then $\frac{\partial k^2 e^{-k(r^2 + r_0^2 e^{-\gamma t})} P(r_0)}{\partial r_0}$ and $\frac{\partial I_1(-2krr_0 e^{-\frac{\gamma t}{2}}) k^2 e^{-k(r^2 + r_0^2 e^{-\gamma t})} P(r_0)}{\partial r_0}$ will also be negative. Hence for $\frac{\partial P(r_0)}{\partial r_0} < 0$, $\frac{\partial P(re^{i\theta}, t)}{\partial r}$ is negative at any time and thus passive.

b) Let us assume $\gamma < 0$. Suppose that the distribution is concentrated about radius r such for long times $\frac{\gamma 2rr_0 e^{-\frac{\gamma t}{2}}}{D(1-e^{-\gamma t})} \rightarrow -\frac{\gamma 2rr_0 e^{-\frac{\gamma t}{2}}}{D} \rightarrow 0$. Then at long times

$$\begin{aligned} P(re^{i\theta}, t) = & \int r_0 dr_0 d\theta_0 \frac{\gamma e^{-\frac{\gamma(r^2 + r_0^2 e^{-\gamma t} - 2rr_0 e^{-\frac{\gamma t}{2}} \cos(\theta - \theta_0))}}{\pi D(1-e^{-\gamma t})} P(r_0 e^{i\theta_0}) \\ \rightarrow & \int r_0 dr_0 d\theta_0 \frac{\gamma e^{-\frac{\gamma(r^2 + r_0^2 e^{-\gamma t})}}{\pi D(1-e^{-\gamma t})} P(r_0 e^{i\theta_0}) = \\ & \frac{\gamma e^{-\frac{\gamma r^2}{D(1-e^{-\gamma t})}}}{\pi D(1-e^{-\gamma t})} \langle e^{-\frac{\gamma r_0^2 e^{-\gamma t}}{D(1-e^{-\gamma t})}} \rangle \end{aligned} \quad (S19)$$

where we have used the fact that

$$\begin{aligned} \int r_0 dr_0 d\theta_0 e^{-\frac{\gamma r_0^2 e^{-\gamma t}}{D(1-e^{-\gamma t})}} P(r_0 e^{i\theta_0}) = \langle e^{-\frac{\gamma r_0^2 e^{-\gamma t}}{D(1-e^{-\gamma t})}} \rangle = \\ \langle e^{-\frac{\gamma n_0 e^{-\gamma t}}{D(1-e^{-\gamma t})}} \rangle, \end{aligned}$$

which is a positive quantity for any distribution. Calculating the derivative over r we obtain at long times

$$\frac{\partial P(re^{i\theta}, t)}{\partial r} = -\frac{2re^{-\frac{\gamma r^2}{D(1-e^{-\gamma t})}}}{(\frac{D}{\gamma}(1-e^{-\gamma t}))^2 \pi} \langle e^{-\frac{\gamma r_0^2 e^{-\gamma t}}{D(1-e^{-\gamma t})}} \rangle < 0 \quad (S20)$$

A counter-example is the coherent state, where the radius of the center of the distribution $r = r_0 e^{-\frac{\gamma t}{2}}$, so

$$\frac{\gamma 2rr_0 e^{-\frac{\gamma t}{2}}}{D(1-e^{-\gamma t})} = \frac{\gamma 2r_0^2 e^{-\gamma t}}{D(1-e^{-\gamma t})} \rightarrow -\frac{\gamma 2rr_0 e^{\frac{\gamma t}{2}}}{D} \rightarrow -\frac{\gamma 2r_0^2}{D}$$

may not be small, and S20 does not hold.

c) It follows from a) and b) that what determines whether or not a distribution becomes passive at long times is the evolution of the center of the distribution, where the derivative vanishes. This is given by

$$\begin{aligned} \frac{\partial P(re^{i\theta}, t)}{\partial r} = & -2 \int dr_0 d\theta_0 (r - r_0 e^{-\gamma t/2} \cos(\theta - \theta_0)) r_0 \times \\ & \frac{e^{-\frac{\gamma |(re^{i\theta} - r_0 e^{-\gamma t/2} e^{i\theta_0})|^2}}{D(1-e^{-\gamma t})}}{\pi (\frac{D}{\gamma}(1-e^{-\gamma t}))^2} P(r_0 e^{i\theta_0}) = 0 \\ \frac{\partial P(re^{i\theta}, t)}{\partial \theta} = & 2 \int dr_0 d\theta_0 (rr_0 e^{-\gamma t/2} \sin(\theta - \theta_0)) r_0 \times \\ & \frac{e^{-\frac{\gamma |(re^{i\theta} - r_0 e^{-\gamma t/2} e^{i\theta_0})|^2}}{D(1-e^{-\gamma t})}}{\pi (\frac{D}{\gamma}(1-e^{-\gamma t}))^2} P(r_0 e^{i\theta_0}) = 0 \end{aligned} \quad (S21)$$